


# Bard Offers Endovascular Stent Grafts with the:


- Broadest Indications in AV Access\*
- Most Complete Size Options\*

**FLAIR®**  
Endovascular Stent Graft




Venous Anastomosis of AV Grafts


**FLUENCY® PLUS**  
Endovascular Stent Graft




AV Fistulae In-Stent Restenosis



Central Veins In-Stent Restenosis\*\*



AV Grafts In-Stent Restenosis



AV Grafts (Venous Outflow)

\* As of June 2016.

\*\* Includes peripheral and central veins (brachiocephalic and subclavian)

The FLAIR® Endovascular Stent Graft is indicated for use in the treatment of stenoses at the venous anastomosis of ePTFE or other synthetic arteriovenous (AV) access grafts.

The FLUENCY® PLUS Endovascular Stent Graft is indicated for use in the treatment of in-stent restenosis in the venous outflow of hemodialysis patients dialyzing by either an arteriovenous (AV) fistula or AV graft and for the treatment of stenosis in the venous outflow of hemodialysis patients dialyzing by an AV graft.

Please consult product labels and instructions for use for indications, contraindications, hazards, additional warnings and precautions.

Bard, Advancing Lives and the Delivery of Health Care, Flair, and Fluency Plus are trademarks and/or registered trademarks of C. R. Bard, Inc. or an affiliate. Copyright © 2017 Bard Peripheral Vascular, Inc. All Rights Reserved. Illustrations by Mike Austin. Copyright © 2017. All Rights Reserved. Bard Peripheral Vascular, Inc. | 1 800 321 4254 | www.bardpv.com | 1625 W. 3rd Street Tempe, AZ 85281 BPV/SGF2/1216/0063